



ELSEVIER

Aquaculture 181 (2000) 257–268

www.elsevier.nl/locate/aqua-online

Aquaculture

Cultivation of the Mediterranean amberjack, *Seriola dumerili* (Risso, 1810), in submerged cages in the Western Mediterranean Sea

Antonio Mazzola, Eugenia Favalaro, Gianluca Sarà *

Laboratory of Marine Biology and Resources, Department of Animal Biology, University of Palermo, Via Archirafi, 18, 90123 Palermo, Italy

Accepted 21 June 1999

Abstract

The growth rate, survival and food conversion ratio (FCR) of the Mediterranean amberjack (*Seriola dumerili*, Risso, 1810) was ascertained in cultivation using submerged net cages in the Gulf of Castellammare (NW Sicily), from September to December 1994. Two net cages (volume = 75 m³) were placed at a depth of 10 m in a sheltered area 1000 m off the coast. Juveniles (mean total length = 141.4 ± 34.2 mm; mean total wet weight = 48 ± 28.1 g) were caught in the gulf under floating wreckage with a purse seine and transplanted to the cages ($n = 800$ per cage) in August. Fish in one cage, group A, were fed with fish scraps whilst fish in group B were fed with pellets. The total length and body wet weight were recorded each month and compared with the wild population of the gulf. Negligible mortality occurred due to capture and transportation to the cages and no diseases were found during the rearing period. Group A reached a final size of 438.1 ± 25.3 mm and 1149 ± 172.2 g, while group B reached 347 ± 25.6 mm and 576 ± 139 g. At this time the wild population was 404.13 ± 17 mm and 777 ± 89.4 g. Food conversion ratios of 1.22 for group A and 3.51 for group B were in the low range compared with other research on *S. dumerili* in the Mediterranean. *S. dumerili* seemed to find fish scraps more appetising than pellets. The low level of investment required, limited breeding period and compatibility with small-scale fishing make submerged net cages a promising system of cultivation for the conversion of coastal fishing. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: *Seriola dumerili*; Cage aquaculture; Open-sea; Feeding; Mediterranean Sea

* Corresponding author. Tel.: +39-91-6167497; fax: +39-91-6172009; E-mail: gsara@mbx.unipa.it



1. Introduction

The Mediterranean amberjack, *Seriola dumerili* (Risso, 1810), is a carnivorous pelagic fish (Carangidae) that is widespread along the Mediterranean coast (Fisher et al., 1987). This species lives at depths of 20–70 m, although it has also been caught at depths down to 360 m (Fisher et al., 1981; Fisher et al., 1987). The young are gregarious by nature and often associated with flotsam, together with *Naucrates ductor* (pilot fish) and *Coryphaena hippurus* (common dolphinfish). *Seriola dumerili* is an opportunistic species, with its diet varying as a function of its size (Mazzola et al., 1993; Matallanas et al., 1995; Pipitone and Andaloro, 1995). It represents an important resource for the southern Mediterranean countries (Andaloro et al., 1992).

In recent years, Mediterranean aquaculture has attempted to select new species of marine fish in order to diversify its production (Cahiers Options Méditerranéennes, 1995). The potential success of a species is based on market analysis, growth performance and on the availability of juveniles. *S. dumerili* is certainly a favourable species for Mediterranean aquaculture to develop. Amberjack are highly adaptable to culture in concrete tanks (Lazzari and Barbera, 1989; Garcia-Gomez, 1993; Greco et al., 1993) and in net cages (Boix et al., 1993; Mazzola et al., 1996), with high growth and survival rates (Porrello et al., 1993; Mazzola et al., 1996).

Research in the Mediterranean area has encountered great difficulty in obtaining spawning in captivity (Lazzari and Barbera, 1989; Manganaro et al., 1993; Lazzari and Di Bitetto, 1994; Marino et al., 1995a,b), despite the success obtained with the Japanese *Seriola quinqueradiata* (Masuma et al., 1990; Tachihara et al., 1993). The provision of 0+ juveniles represents a limit in the development of the cultivation of *S. dumerili* (Greco et al., 1991; Caridi et al., 1992; Mazzola et al., 1996). One solution to this problem could be the provision of juveniles in their natural habitat, although overfishing has caused a reduction in this resource (Mazzola et al., 1993).

Little information is available regarding the biology of the Mediterranean amberjack in cultivation (Giovanardi et al., 1984; Navarro et al., 1987; Porrello et al., 1993), its capture, transport (Greco et al., 1991) and dietary preferences (Garcia-Gomez, 1993).

This paper presents the results of the cultivation of Mediterranean amberjack in submerged cages using simple installation techniques of low technology and cost. Research was carried out to test the growth rate, survival and feeding parameters in captivity of two groups of fish fed two different diets. Their growth rates were compared with that of the wild population taken from the same area.

2. Materials and methods

Feeding trials were carried out between July 1994 and December 1994 in the Gulf of Castellammare (NW Sicily; 38°02'31"N; 12°55'28"E) in a sheltered area 1000 m off the coast. Collection of the juvenile fish and management of the cultivation activities were carried out by a co-operative of local fishermen.



Two cylindrical net cages of 75 m³ (ϕ 4.5 m, h 5 m, and 12-mm mesh size) were moored onto artificial reefs, 18 m deep, and submerged 10 m below the water surface. Feeding was carried out by means of a mesh cone from the cage to the surface.

Fishermen caught the juveniles during the summer of 1994 from below fish aggregating devices (FAD). These FADs (artificial wreckage made of twisted cane mats) were




Fig. 1. Monthly trends of total length (a) and wet weight (b) of *Seriola dumerili* in cage culture and the wild.

moored to the sea bottom along a bathymetric gradient (5–120 m). The juveniles were caught with a small purse seine (150 m × 40 m and 20-mm mesh) and the total length (TL) and wet weight (T_{ww}) of sampled specimens recorded immediately. The fish were placed in PVC tanks (volume: 0.3 m³) on the fishing boat, which was equipped with an open forced seawater system for transplanting to the cages (within 4 h). Amberjack specimens were placed into the cages ($n = 800$ per cage) at an initial density of 10 specimens m⁻³.

During the rearing period (September–December), monthly samples of *Seriola dumerili* were taken ($n = 50$ per cage), TL and T_{ww} were measured and the fish then replaced inside the cages. At the same time, samples of wild *S. dumerili* were caught by the same group of fishermen in the Gulf of Castellammare and treated as above.

The first group of fish (A) to be placed in cultivation was fed with fish scraps obtained from Sicilian fisheries, while the second group (B) was fed with pellets (Trouvit Marine; Hendrix). The two groups were fed twice a day (at 7 am and 7 pm) and the total amount of food administered was determined according to the classic methods used in Mediterranean aquaculture (Lazzari and Barbera, 1989).

Growth performance was analysed by calculating the daily specific growth rate (Winberg, 1971) in somatic wet weight (C_w) and in length (C_l) ($C_{w/l} = [\ln X_2 - \ln X_1] / \Delta t$) where X represented length (TL) or weight (W). The relationship between the total length and somatic wet weight was calculated using a simple allometric equation ($W = aL^b$; Gould, 1966) and logarithmic transformations. The relative condition factor (K_n , Le Cren, 1951) was also calculated by using the follow relationship: $K_n = W_{(meas)} / [aL^b]$ where $W_{(meas)}$ represented our measured weights and $[aL^b]$ represented weight estimated by means allometric equations. Feeding was studied by calculating the daily rate of feeding (f ; Tsevis et al., 1992) according the equation:




Fig. 2. Monthly trends of relative condition factor (K_n ; Le Cren, 1951) of *Seriola dumerili* in cage culture and the wild, relative to water temperature changes during cultivation.

$f = F_t * 100 / [t * (W_0 + W_t + W_d) / 2]$, where F_t was the total food consumption, W_0 and W_t were respectively the initial and final biomass, W_d the dead biomass and t the time in days and the food conversion ratio (FCR; Carter et al., 1994) according the follow equation: $FCR = FT_{DW} / (W_0 - W_t)$ where FT_{DW} was total dry food and W_0 and W_t were the same as above. The Kruskal–Wallis test (Sokal and Rohlf, 1981) was used to ascertain differences in growth performance.




Fig. 3. Specific daily growth rates in length (a) and wet weight (b) relative to the temperature trend for the 3 groups of fish in cultivation.

3. Results

Water temperature at the surface ranged from 16.7°C (autumn–winter) to 22°C (summer) (mean value = $19.58 \pm 2.03^\circ\text{C}$) and salinity was between 37.5 psu (autumn–winter) and 38.1 psu (summer) (mean value = 37.88 ± 0.19 psu). The thermocline was observed to be at a depth of approximately 10–12 m for the whole summer period until September.

3.1. Cultivation phase

3.1.1. Growth

Fig. 1a and b show the monthly trend of total length and total wet weight of the 3 groups of *Seriola dumerili* during the experimentation period. Although there were no significant differences in wet weight ($H_{\text{TWw}} = 6.93$; $P > 0.05$; $n = 12$), group A reached the largest size. Significant differences were, however, detected in total length ($H_{\text{TL}} = 7.93$; $P < 0.05$; $n = 12$). The overlapping of the growth curves for the wild population and group B was also confirmed by the relative condition factor (K_n) trend (Fig. 2).

The relative condition factor showed that increases in weight and length did not lead to a corresponding increase in this factor: when the weight increased from 158 g to 777 g (October, group A), the condition factor decreased from 1.04 to 1.01. These values were lower than those of the wild population. These group A specimens increased a great deal in size and represented a departure from the general trend shown by the wild population.




Fig. 4. Monthly trend of allometric coefficient ($b = \text{slope of allometric regression}$) of fish in cage and wild in comparison with a typical trend as indicated in Gould (1966).

Table 1
Diet composition and biochemical features of the main species composing fish scrap supplied to the *Seriola dumerili* of group A. Biochemical composition of food derived by Bono et al., 1992

| Species | % Composition in diet | Total amount of food in dry weight (kg) | % Protein | % Lipid | % Carbohydrate | % Ash |
|-------------------------------|-----------------------|---|-----------|---------|----------------|-------|
| <i>Scomber scomber</i> | 46.38 | 496.5 | 48.51 | 31.67 | 3.99 | 15.83 |
| <i>Boops boops</i> | 16.93 | 157.3 | 59.66 | 18.5 | 7.36 | 14.48 |
| <i>Trachurus trachurus</i> | 15.51 | 155.9 | 73.44 | 4.16 | 1.24 | 21.16 |
| <i>Spicara</i> sp. | 3.865 | 29.54 | 68.81 | 2.91 | 0.67 | 27.61 |
| <i>Lepidotrigla cavillone</i> | 2.68 | 19.63 | 81.19 | 4.93 | 0.96 | 12.92 |
| <i>Merluccius merluccius</i> | 2.623 | 18.27 | 85.13 | 3.81 | 0.64 | 10.42 |
| <i>Lophius budegassa</i> | 1.128 | 8.316 | 69.63 | 2.68 | 0.77 | 26.92 |
| <i>Arnoglossus laterna</i> | 0.31 | 2.349 | 67.88 | 8.97 | 0.98 | 22.17 |
| <i>Lepidopus caudatus</i> | 0.282 | 2.464 | 65.38 | 13.53 | 1.12 | 19.97 |
| <i>Argentina sphyraena</i> | 0.141 | 1.106 | 75.69 | 13.06 | 0.74 | 10.51 |
| <i>Capros aper</i> | 0.141 | 1.241 | 61.56 | 16.84 | 1.16 | 20.44 |
| <i>Gnathophipis mystax</i> | 0.085 | 0.648 | 80.19 | 3.65 | 0.94 | 15.22 |
| Other species | 9.929 | 74.43 | 69.76 | 10.39 | 1.71 | 18.14 |
| Total amount of food | 100 | 967.6 | — | — | — | — |

Table 2

Diet composition and biochemical features as reported by Hendrix (Italy) of pelleted food supplied to *Seriola dumerili* of group B

| Composition | Pellet size (mm) | | |
|--------------------|------------------|------|------|
| | 3.00 | 4.00 | 8.00 |
| % H ₂ O | 8.20 | 9.00 | 10.0 |
| % Protein | 47.0 | 47.5 | 48.0 |
| % Lipid | 18.0 | 18.0 | 10.5 |
| % Cellulose | 1.30 | 1.00 | 1.00 |
| % Ash | 12.6 | 11.2 | 11.0 |

No significant differences were shown in specific growth rate between the three groups ($H_{CL} = 5.62$; $P > 0.05$; $n = 12$; $H_{CWW} = 4.85$; $P > 0.05$; $n = 12$) (Fig. 3a, b), even though group A showed a peak in September–October and November–December. The two other populations showed a constant trend, except for the months of November–December when the rate reached minimum value.

The coefficient b trend of the allometric relationship between TL and T_{WW} (Fig. 4) as an index of growth performance (Gould, 1966) displayed the typical trend for the wild population at around a mean value of 2.80 ± 0.26 . Group A gave similar results with a mean value of 2.82 ± 0.71 except in November when b reached an anomalous value of 4, while group B showed a different trend with higher values (mean = 3 ± 0.33).

3.1.2. Feeding features

Tables 1 and 2 (data according to Bono et al., 1992 and Trouvit, respectively) report the diet and chemical composition of the food used in cultivation, while Table 3a–b show the feeding features for the two groups of *S. dumerili*. The total amount of food supplied to group A was 3545 kg (967.6 kg dry matter) during the period of cultivation. The diet was principally composed of *Scomber scomber* (46.38%), *Boops boops*

Table 3

(a) Feeding parameters of the *Seriola dumerili* of group A

| Month | Total wet biomass (kg) | Total amount dry weight of food (kg) | Food supplied as a function of body weight (%) | Density (kg m ⁻³) | Daily rate of feeding |
|-------|------------------------|--------------------------------------|--|-------------------------------|-----------------------|
| SEPT | 126.40 | 93.14 | 9.00 | 1.68 | 3.01 |
| OCT | 621.60 | 267.17 | 5.25 | 8.28 | 1.90 |
| NOV | 619.20 | 261.07 | 5.15 | 8.26 | 1.12 |
| DEC | 919.20 | 346.17 | 4.60 | 12.30 | 1.20 |

b. Feeding parameters of the *Seriola dumerili* of group B

| | | | | | |
|------|--------|--------|------|------|------|
| SEPT | 108.00 | 117.86 | 4.00 | 1.44 | 4.29 |
| OCT | 224.00 | 244.45 | 4.00 | 2.99 | 3.93 |
| NOV | 456.00 | 373.22 | 3.00 | 6.08 | 2.93 |
| DEC | 460.80 | 502.86 | 4.00 | 6.14 | 2.93 |

(16.93%) and *Trachurus trachurus* (15.51%), with these three species representing over 78% of the total amount of food consumed. Group A reached a total wet biomass of 919.2 kg, a final density in the cage of 12.3 kg/m³ and a food conversion ratio of 1.22. The daily feeding rate was higher during September (3.01% T_{ww}), decreasing to 1.20% T_{ww} in December.

The total amount of food supplied to group B was 1362 kg (1,238 kg dry matter). Group B fish reached 461 kg in total biomass, a final density of 6.14 kg/m³ and a food conversion ratio of 3.51. The daily feeding rate was found to be more constant during the cultivation period, dropping slightly from September (4.29% T_{ww}) to December (2.93% T_{ww}).

4. Discussion

In this study the survival rate of captured juveniles was 100%, compared with 98% reported by Porrello et al. (1993) and 80% by Cavaliere et al. (1989). Mortality ($n = 25$; 3.12%) was then observed only at the beginning of the experiment (in September) in the pellet-fed group. This was probably due to acclimatization and not to the cultivation process. Temperature changes were not observed to have a negative influence, contrary to the findings of Garcia-Gomez (1993).

The two diet types did not significantly affect the growth trends, although group A reached a larger size than the other two groups. Similar findings were observed in *Seriola quinqueradiata* fed both with different pellet types (Shimeno et al., 1993) and with raw fish (Watanabe et al., 1993).

Specific daily growth rate, the condition factor and the allometric relationship did not show a correlation with temperature because the cages were located beneath the thermocline where only small variations in temperature occurred ($\Delta T + 4^{\circ}\text{C}$). In contrast, wide monthly changes in chemical–physical variables have been shown to have an important role in growth in floating cages and tanks (Garcia-Gomez, 1993; Porrello et al., 1993).

The results of other experiments carried out with *Seriola dumerili* in the western Mediterranean area are reported in Table 4. Different results were obtained from experiments in tanks by Garcia-Gomez (1993). Growth curves of pellet-fed *S. dumerili* were similar to those of the wild population, although the maximum size was not significantly different.

The growth trend and the maximum size of group A were better than those obtained in floating cages (Giovanardi et al., 1984; Porrello et al., 1993) and tanks (Cavaliere et al., 1989; Garcia-Gomez, 1993; Greco et al., 1993). The food conversion ratio obtained with pellets was high despite their elevated nutritional value (Garcia-Gomez, 1993) and higher compared to the ratio of group A. Group A was fed with fish scraps which, despite their high water content, were found to be more nutritious. Food scraps produced a food conversion ratio that was lower than those reported in the literature.

The greatest degree of homogeneity in the class sizes was observed in group B (Garcia-Gomez, 1993), and this was probably due to the homogeneous size of the pellets compared to the fish scraps, which varied in weight and shape. Furthermore, the fish



Table 4
Comparing results of other experiments of cultivation of *Seriola dumerili* carried out in the Mediterranean

| Site | Type of cultivation | Initial total wet weight (g) | Final total wet weight (g) | Time of cultivation (day) | Diet type | Food conversion ratio | References |
|--------------------------|---------------------|------------------------------|----------------------------|---------------------------|-----------------------------|-----------------------|-------------------------|
| Castellammare (Sicily) | Floating cage | 130 | 670 | 85 | Fish scraps | Not reported | Giovanardi et al., 1984 |
| Messina (Sicily) | PVC tanks | < 100 (?) | 1214 | 380 | Fish scraps | Not reported | Cavaliere et al., 1989 |
| Messina (Sicily) | PVC tanks | 58.1 | 379 | 110 | Fish scraps | 3 | Cavaliere et al., 1989 |
| Coasts of Spain | PVC tanks | 64 | 412 | 120 | Pellets | 1.80 | Garcia-Gomez, 1993 |
| Coasts of Spain | PVC tanks | 64 | 367 | 120 | Pellets | 2.30 | Garcia-Gomez, 1993 |
| Coasts of Spain | PVC tanks | 64 | 396 | 120 | Fish scraps | 4.89 | Garcia-Gomez, 1993 |
| Coasts of Spain | PVC tanks | 64 | 342 | 90 | Fish scraps | 1.48 | Garcia-Gomez, 1993 |
| Messina (Sicily) | PVC tanks | 106 | 500 | 300 | Fish scraps | 4 | Greco et al., 1993 |
| Messina (Sicily) | PVC tanks | 106 | 392 | 300 | Fish scraps + chicken liver | 5 | Greco et al., 1993 |
| Aeolian Islands (Sicily) | Floating cage | 72.7 | 858 | 120 | Fish scraps | 4.4 | Porrello et al., 1993 |
| Castellammare (Sicily) | Submerged cage | 48 | 576 | 120 | Pellets | 3.51 | This paper |
| Castellammare (Sicily) | Submerged cage | 48 | 1149 | 120 | Fish scraps | 1.22 | This paper |



scraps may have produced competitive behaviour and a degree of territoriality among the specimens, resulting in a marked heterogeneity in class sizes.

This study has shown that cages are easy to use and are compatible with the work of small-scale fishermen. The low investment costs, limited breeding period and suitability for small-scale fishing all make this a useful way to convert coastal fishing. The absence of a visual impact from these cages also makes them an acceptable proposal for sea parks and protected areas.

Acknowledgements

The authors would like to thank the three anonymous referees for their invaluable help in improving the manuscript and Helen Main for her proof-reading. This work was funded by the Ministero Università Ricerca Scientifica e Tecnologica (MURST, Italy) and by the Ministero Politiche Agricole (MiPA, Italy).

References

- Andaloro, F., Potoschi, A., Porrello, S., 1992. Contribution to the knowledge of growth of greater amberjack, *Seriola dumerili* (Cuv., 1817) in the Sicilian Channel (Mediterranean Sea). Rapp. Comm. Int. Mer Médit. 33, 282.
- Boix, J., Fernandez, J.V., Macia, G., 1993. El cultivo de *Seriola* (*Seriola dumerili*, Risso, 1810) en jaula flotante en el termino marítimo de Les Cases d'Alcanar (Tarragona). Actas IV Congreso Nacional Acuicultura. La Toja, Galicia, 133–136 pp.
- Bono, G., Arculeo, M., Mazzola, A., Greco, S., 1992. Lo scarto di pesca come risorsa utilizzabile in acquicoltura. Oebalia 17 (2 suppl.), 71–76.
- Cahiers Options Méditerranéennes, 1995. Marine aquaculture finfish species diversification. CIHEAM/MEDRAP II, 16: 1–169.
- Caridi, D., Greco, S., Lo Paro, G., Micale, V., Arena, G., 1992. Indagini di pesca per il reperimento di novellame di *Seriola dumerili* in un'area del basso Tirreno (Is. Eolie). Oebalia 17, 91–96.
- Carter, C.G., Houlihan, D.F., Buchanan, B., Mitchell, A.I., 1994. Growth and feed utilization efficiencies of sea water Atlantic salmon, *Salmo salar* L., fed a diet containing supplementary enzymes. Aquacult. Fish. Manage. 25, 37–46.
- Cavaliere, A., Crisafi, E., Faranda, F., Greco, S., Lo Paro, G., Manganaro, A., Mazzola, A., 1989. Collection of fingerlings and rearing of *Seriola dumerili* in tanks. In: De Pauw, N., Jaspers, E., Ackefors, H., Wilkins, N. (Eds.), Aquaculture A Biotechnology in progress. Bredene, Belgium. EAS: 119–123 pp.
- Fisher, W., Bianchi, G., Scott W.B., 1981. FAO specie identification sheets for fishery purposes. Eastern Central Atlantic (fishing area 34, 47 (in part)). Canad. Funds in Trust. Ottawa, Department of Fisheries and Oceans Canada, FAO of the U.N., 3.
- Fisher, W., Bauchot, M.L., Schneider, M., 1987. Fiche FAO d'identification des especes pour les besoins de la peche. (Rev. 1). Méditerranée et Mer Noire (Zone de peche 37). FAO Project GCP/INT/422/EEC, 1529 pp.
- García-Gomez, A., 1993. Primeras experiencias de crecimiento de juveniles de *Seriola mediterranea* (*Seriola dumerili*, Risso, 1810) alimentados con una dieta semihumeda. Bol. Inst. Esp. Oceanogr. 9 (2), 347–360.
- Giovanardi, O., Mattioli, G., Piccinetti, C., Sambucci, G., 1984. Prime esperienze sull'allevamento di *Seriola dumerili* (Risso, 1810) in Italia. Riv. Ital. Pisc. Ittiopat. 4, 123–130.
- Gould, S.J., 1966. Allometry and size in ontogeny and phylogeny. Biol. Rev. 41, 587–640.
- Greco, S., Arena, G., Caridi, D., Micale, V., 1991. An improved method of capture and transport for juveniles of *Seriola dumerili*. In: De Pauw, N., Joyce, J. (Eds.), Aquaculture and Environment. Dublin Aquaculture 1991, 14, 130–131.

- Greco, S., Caridi, D., Cammaroto, S., Genovese, L., 1993. Preliminary studies on artificial feeding of amberjack fingerlings. In: Barnabè, G., Kestemont, P. (Eds.), Production, Environment and Quality. Bordeaux Aquaculture 1992. EAS, Special publication Ghent, Belgium 1993, 18, 247–254.
- Lazzari, A., Barbera, G., 1989. Farming the Mediterranean yellowtail, *Seriola dumerili* (Risso, 1810) in concrete ponds: results and perspectives. In: De Pauw, N., Jaspers, E., Ackefors, H., Wilkins, N. (Eds.), Aquaculture A Biotechnology in progress. Bredene, Belgium. EAS: 209–213.
- Lazzari, A., Di Bitetto, M., 1994. Maturity stages determination in *Seriola dumerili* using ovary histology techniques. Bordeaux, Aquaculture 1994, 25–33 pp.
- Le Cren, E.D., 1951. The length–weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). J. Anim. Ecol. 20, 201–219.
- Manganaro, A., Barbera, G., Cammaroto, S., Greco, S., 1993. Campagna di cattura di *Seriola dumerili* e tentativi di riproduzione artificiale. Biol. Mar. Suppl. Not. Italian Marine Biology Society 1, 245–249.
- Marino, G., Mandich, A., Massari, A., Andaloro, F., Porrello, S., Finoia, M.G., Cevalco, F., 1995a. Aspects of reproductive biology of the Mediterranean amberjack (*Seriola dumerili* Risso, 1810) during the spawning period. J. Appl. Ichthyol. 11, 9–24.
- Marino, G., Porrello, S., Andaloro, F., Massari, A., Mandich, A., 1995b. Aspects of reproductive biology of the Mediterranean amberjack (*Seriola dumerili* Risso, 1810): gonadal development. Cah. Options Méditerran. 16, 115–124.
- Masuma, S., Kanematu, M., Teruya, K., 1990. Embryonic and morphological development of larvae and juveniles of the amberjack, *Seriola dumerili*. Jpn. J. Ichthyol. 37, 164–169.
- Matallanas, J., Casadevall, M., Carrasson, M., Boix, J., Fernandez, V., 1995. The food of *Seriola dumerili* (Pisces: Carangidae) in the Catalan sea (Western Mediterranean). J. Mar. Biol. Assoc. U.K. 75, 257–260.
- Mazzola, A., Lopiano, L., Sarà, G., D'Anna, G., 1993. Sistemi di pesca, cattura ed abitudini alimentari di *Seriola dumerili* (Risso, 1810) nel Golfo di Castellammare (Sicilia Occidentale) (Pisces: Perciformes). Naturalista Siciliano 12 (1–2), 137–148.
- Mazzola, A., Sarà, G., Favaloro, E., Mirto, S., 1996. Sistemi di maricoltura open-sea per l'allevamento di *Seriola dumerili* (Pisces, Osteichthyes) nel Golfo di Castellammare (Sicilia Occidentale). Biol. Mar. Medit. 3 (1), 176–185.
- Navarro, J.M., Belmonte, A., Culmarex, S.a.L., 1987. Cultivo de *Seriola* en jaulas flotantes en la bahia de El Hornillo (Murcia SE Espana). Cuad. Marisq. Publ. Tecn. 12, 11–16.
- Pipitone, C., Andaloro, F., 1995. Food and feeding habits of juvenile greater amberjack, *Seriola dumerili* (Osteichthyes, Carangidae) in inshore waters of the central Mediterranean Sea. Cybium 19 (3), 305–310.
- Porrello, S., Andaloro, F., Vivona, P., Marino, G., 1993. Rearing trial of *Seriola dumerili* in a floating cage. In: Barnabè, G., Kestemont, P. (Eds.), Production, Environment and Quality. Bordeaux Aquaculture 1992. EAS, Special publication Ghent, Belgium 1993, 18, 299–307.
- Shimeno, S., Kumon, M., Ando, H., Ukawa, M., 1993. The growth performance and body composition of young yellowtail fed with diets containing defatted soybean meal for a long period. Nippon Suisan Gakkaishi Bull. Jap. Soc. Sci. Fish. 59 (5), 821–825.
- Sokal, R.R., Rohlf, F.J. Biometry. W.H. Freeman, New York, 1981, pp. 859.
- Tachihara, K., Ebisu, R., Tukahima, Y., 1993. Spawning, eggs, larvae and juvenile of the purplish amberjack *Seriola dumerili*. Nippon Suisan Gakkaishi 59 (9), 1479–1488.
- Tsevis, N., Klaoudatos, S., Conides, A., 1992. Food conversion budget in sea bass *Dicentrarchus labrax*, fingerlings under two different feeding frequency patterns. Aquaculture 101, 293–304.
- Watanabe, T., Takeuchi, T., Okamoto, N., Viyakarn, V., Sakamoto, T., Satoh, S., Matsuda, M., 1993. Feeding experiments of yellowtail with a newly developed soft-dry pellet. J. Tokyo Univ. Fish. 80 (1), 1–17.
- Winberg, G.G. Methods for the estimation of production of aquatic animals. Academic Press, London, 1971, 175 pp.

